



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## CALCULUS.

—

263. Proposed by V. M. SPUNAR, M. S., C. E., East Pittsburg, Pa.

Find a point such that the sum of the squares of its distances from  $n$  given points shall be a minimum, and prove that the value so found is  $1/n$ th part of the sum of the squares of the mutual distances between the  $n$  points, taken two and two.

264. Proposed by W. J. GREENSTREET, M. A., Stroud, England.

The join of the center of curvature of a curve to the origin is at  $\alpha$  to the initial line. Prove that with the usual notation:

$$\frac{d\alpha}{d\psi} \left[ \left( \frac{dp}{d\psi} \right)^2 + \left( \frac{d^2p}{d^2\psi} \right)^2 \right] = \frac{dp}{d\psi} \frac{d\rho}{d\psi}.$$

## MECHANICS.

—

219. Proposed by W. J. GREENSTREET, M. A., Stroud, England.

A rod length  $a\sqrt{3}$ , weight  $W$ , has at each end a smooth ring which can slide on a vertical circle radius  $r$ . Each ring is attached by an elastic string (natural lengths  $a, b$ ; moduli  $\mu a, \mu b$ ) to the highest point of the circle. Find the inclination of the rod to the horizon in a position of equilibrium.

220. Proposed by W. J. GREENSTREET, M. A., Stroud, England.

Four particles  $A, B, C, D$  lie on a smooth table at the corners of a rhombus.  $AB, BC, CD, DA$  are light inextensible strings connecting the particles. The angle at  $A$  is acute. A blow is given to  $A$  along the diagonal, away from  $C$ . Find the ratio of the initial velocity of  $C$  to that of  $A$ .

## NUMBER THEORY AND DIOPHANTINE ANALYSIS.

—

155. Proposed by PROF. R. D. CARMICHAEL, Anniston, Alabama.

If  $p$  and  $q$  are primes and  $m$  and  $n$  are any integers, find the cases in which the equation  $p^m - q^n = 1$  may be satisfied.

156. Proposed by A. H. HOLMES, Brunswick, Maine.

Find integral values for  $a, b, c, d$ , and  $e$  in the equation,  $a^2 + b^2 + c^2 + d^2 = e^2$ .

## AVERAGE AND PROBABILITY.

—

199. Proposed by PROF. R. D. CARMICHAEL, Anniston, Ala.

A circle is inscribed in a given square. Two points are taken at random within the square but without the circle. What is the chance the distance between the points does not exceed the side of the square?